



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2015

Analgesia and unwanted benzodiazepine effects in point-mutated mice expressing only one benzodiazepine-sensitive GABAA receptor subtype

Ralvenius, William T ; Benke, Dietmar ; Acuña, Mario A ; Rudolph, Uwe ; Zeilhofer, Hanns Ulrich

Abstract: Agonists at the benzodiazepine-binding site of GABAA receptors (BDZs) enhance synaptic inhibition through four subtypes (1, 2, 3 and 5) of GABAA receptors (GABAAR). When applied to the spinal cord, they alleviate pathological pain; however, insufficient efficacy after systemic administration and undesired effects preclude their use in routine pain therapy. Previous work suggested that subtype-selective drugs might allow separating desired antihyperalgesia from unwanted effects, but the lack of selective agents has hitherto prevented systematic analyses. Here we use four lines of triple GABAAR point-mutated mice, which express only one benzodiazepine-sensitive GABAAR subtype at a time, to show that targeting only 2GABAARs achieves strong antihyperalgesia and reduced side effects (that is, no sedation, motor impairment and tolerance development). Additional pharmacokinetic and pharmacodynamic analyses in these mice explain why clinically relevant antihyperalgesia cannot be achieved with nonselective BDZs. These findings should foster the development of innovative subtype-selective BDZs for novel indications such as chronic pain.

DOI: <https://doi.org/10.1038/ncomms7803>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-110282>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Ralvenius, William T; Benke, Dietmar; Acuña, Mario A; Rudolph, Uwe; Zeilhofer, Hanns Ulrich (2015). Analgesia and unwanted benzodiazepine effects in point-mutated mice expressing only one benzodiazepine-sensitive GABAA receptor subtype. *Nature Communications*, 6(6803):online.

DOI: <https://doi.org/10.1038/ncomms7803>

ARTICLE

Received 8 Aug 2014 | Accepted 2 Mar 2015 | Published 13 Apr 2015

DOI: 10.1038/ncomms7803

OPEN

Analgesia and unwanted benzodiazepine effects in point-mutated mice expressing only one benzodiazepine-sensitive GABA_A receptor subtype

William T. Ralvenius^{1,2}, Dietmar Benke^{1,2}, Mario A. Acuña^{1,2}, Uwe Rudolph^{3,4} & Hanns Ulrich Zeilhofer^{1,2,5}

Agonists at the benzodiazepine-binding site of GABA_A receptors (BDZs) enhance synaptic inhibition through four subtypes ($\alpha 1$, $\alpha 2$, $\alpha 3$ and $\alpha 5$) of GABA_A receptors (GABA_AR). When applied to the spinal cord, they alleviate pathological pain; however, insufficient efficacy after systemic administration and undesired effects preclude their use in routine pain therapy. Previous work suggested that subtype-selective drugs might allow separating desired antihyperalgesia from unwanted effects, but the lack of selective agents has hitherto prevented systematic analyses. Here we use four lines of triple GABA_AR point-mutated mice, which express only one benzodiazepine-sensitive GABA_AR subtype at a time, to show that targeting only $\alpha 2$ GABA_ARs achieves strong antihyperalgesia and reduced side effects (that is, no sedation, motor impairment and tolerance development). Additional pharmacokinetic and pharmacodynamic analyses in these mice explain why clinically relevant antihyperalgesia cannot be achieved with nonselective BDZs. These findings should foster the development of innovative subtype-selective BDZs for novel indications such as chronic pain.

¹Institute of Pharmacology and Toxicology, University of Zurich, Winterthurerstrasse 190, CH-8057 Zurich, Switzerland. ²Center for Neuroscience Zurich (ZNZ), Winterthurerstrasse 190, CH-8057 Zurich, Switzerland. ³Laboratory of Genetic Neuropharmacology, McLean Hospital, 115 Mill Street, Belmont, Massachusetts 02478, USA. ⁴Department of Psychiatry, Harvard Medical School, 401 Park Drive, Boston, Massachusetts 02215, USA. ⁵Institute of Pharmaceutical Sciences, Swiss Federal Institute of Technology (ETH) Zurich, Vladimir-Prelog-Weg 4, CH-8093 Zurich, Switzerland. Correspondence and requests for materials should be addressed to H.U.Z. (email: zeilhofer@pharma.uzh.ch).

Chronic pain is a severe medical condition affecting hundreds of millions of patients worldwide. It is widely accepted that diminished inhibition in pain-processing circuits of the spinal dorsal horn is a major contributor to different chronic pain forms^{1–4}. We have previously demonstrated that local spinal application of BDZ site ligands that positively modulate GABA_AR function alleviates neuropathic and inflammatory pain in rodents⁵. Translation of these results to routine systemic pain treatment, however, requires a separation of desired antihyperalgesia from unwanted side effects. This separation appears potentially feasible based on the existence of different GABA_AR subtypes.

Most GABA_ARs in the brain and spinal cord are heteropentameric ion channels composed of two α , two β and one γ 2 subunit⁶. The high-affinity binding site for BDZs is formed by an interface between one α subunit and the γ 2 subunit. High-affinity binding of BDZs at this site requires the presence of a histidine residue at a conserved site in the N-terminal domain of the α subunit. This conserved histidine is present in α 1, α 2, α 3 or α 5 subunits, but not in the α 4 and α 6 subunits⁷. Mutation of the histidine residue into an arginine dramatically reduces the affinity of GABA_ARs to BDZs without changing their responses to GABA⁸. The generation of histidine to arginine (H→R) point-mutated mice for each of the four BDZ-sensitive GABA_AR subunits has allowed attributing the different *in vivo* effects of BDZs to defined GABA_AR subtypes⁹. Most importantly, it was shown that the sedative effects of BDZs were strongly reduced in mice carrying the H→R point mutation in their α 1 subunits^{10,11}, while point-mutating the α 2 subunits led to a loss in the anxiolytic effects of BDZs¹². Using this approach, we could demonstrate that mice whose α 2GABA_ARs had been rendered BDZ-insensitive show drastically reduced antihyperalgesic responses to spinal diazepam (DZP)⁵. A general consensus on the question, which GABA_AR subtype should best be targeted to achieve maximal antihyperalgesic responses and to best avoid undesired effects has not been reached. A similarly open question is why classical BDZs are largely devoid of analgesic actions in patients.

Highly selective tool compounds that would allow addressing these questions pharmacologically are still lacking¹³. For this reason, we have generated triple GABA_AR point-mutated mice, in which only a single GABA_AR subtype remains BDZ-sensitive. We designate these mice hereafter as HRRR, short for α 1H; α 2R; α 3R; α 5R (that is, a mouse in which only α 1GABA_ARs are left BDZ-sensitive) and, accordingly, as RHRR, RRHR, RRRH for mouse lines, in which either only α 2, α 3 or α 5 subunits remain BDZ-sensitive. In these mice, the action of the classical normally nonselective BDZ agonist DZP should mimic the action of fully subtype-selective compounds. Such a genetic approach also avoids possible confounding factors of subtype-selective compounds such as pharmacokinetic differences and unknown specificity profiles of drug metabolites. With the use of triple GABA_AR point-mutated mice, we were not only able to perform a systematic analysis on the GABA_AR subtypes best targeted for optimal antihyperalgesia and least-pronounced side effects; however, we also provide an explanation why classical (nonselective) BDZs are largely devoid of clinically relevant analgesic properties.

Results

Spinal distribution of subtype-specific BDZ-binding sites. In the first series of experiments, we characterized the distribution pattern of the four subtypes of BDZ-sensitive GABA_ARs ('BDZ receptors') in the spinal dorsal horn of triple point-mutated mice. Using immunocytochemistry, we verified that neither the regional distribution nor the expression levels of α 1, α 2, α 3 and α 5

subunits and of the γ 2 subunit differed between wild-type (wt) mice and the four strains of triple point-mutated mice (Fig. 1a,b). Similarly, the total number of BDZ-binding sites (wt receptors plus H→R point-mutated receptors) quantified through [³H]Ro15-4513 autoradiography was unchanged (Fig. 1c). Spinal autoradiography with [³H]flumazenil that binds with high affinity only to nonpoint-mutated GABA_ARs allowed a quantitative analysis of each of the four subtypes of BDZ-binding sites in isolation. In wt mice, [³H]flumazenil exhibited specific binding throughout the spinal grey matter with enrichment in the dorsal horn and around the central canal. The density of binding sites in the dorsal horn was highest for the α 3GABA_AR subtype, followed by α 2 and α 1, and lowest for the α 5 subtype (Fig. 1d). The α 1 subtype was concentrated around the central canal, α 2 was most abundant in the superficial dorsal horn, where nociceptive fibres terminate, and α 3 was found throughout the dorsal horn and around the central canal, while α 5 was generally weak. The distribution of subtype-specific [³H]flumazenil binding in the four strains of GABA_AR triple point-mutated mice matched the distribution of the subunits determined with immunocytochemistry on a gross scale. However, the sum of the binding sites in the four strains of triple point-mutated mice was about twice as high as the total number of binding sites in wt mice. The results from the immunocytochemistry and [³H]Ro15-4513 autoradiography largely rule out changes in protein expression as the underlying cause. Instead, the [³H]flumazenil data are consistent with a high prevalence of GABA_ARs containing two different α subunits^{14,15} and with a model of the GABA_AR assembly in which nonpoint-mutated α subunits have a higher probability for interaction with the γ 2 subunit than H→R point-mutated subunits¹⁶. This conclusion is further supported by the only marginal reduction of BDZ binding observed in spinal cords of the four single point-mutated mouse lines (Fig. 1e).

Antihyperalgesia by single BDZ-sensitive GABA_ARs subtypes.

We next used the triple point-mutated mice to predict pharmacological actions of subtype-selective BDZ site agonists and treated them with DZP, a classical nonselective BDZ, whose agonistic activity was restricted to a single GABA_AR subtype in triple point-mutated mice. We first focused on the antihyperalgesic potential of such an approach (that is, the potential to reduce neuropathy-induced hyperalgesia) and tested DZP in mice subjected to the chronic constriction injury (CCI; ref. 17) model. Baseline mechanical and heat sensitivity were statistically indistinguishable in all strains of mice analysed and all strains of mice developed similar hyperalgesia after CCI (Table 1). All subsequent experiments with DZP were performed on an α 1R point-mutated background to avoid DZP-induced sedation, which is a potential confounding factor in behavioural pain tests. We first tested different doses of systemic (per os (p.o.)) DZP in α 1 point-mutated RHHH mice (Fig. 2a,b). On the basis of these results, we chose a dose of 10 mg kg^{−1} body weight, which is close to the ED₇₀, for subsequent experiments. We then asked which GABA_AR subtype exerts the strongest antihyperalgesic action and tested the effects of DZP on mechanical (von Frey filament test) and heat (Hargreaves test) hyperalgesia in the four strains of GABA_AR triple point-mutated mice (Fig. 2c–e). For both stimuli, the strongest effect was achieved in RHRR mice in which only α 2GABA_ARs were BDZ-sensitive. Exclusive targeting of α 3 and α 5GABA_ARs (in RRHR and RRRH mice) also elicited antihyperalgesia but to a smaller extent. Importantly, mice in which all four BDZ receptors had been point-mutated (RRRR mice) did not show any antihyperalgesic response. We next asked whether targeting of a second or third GABA_AR subtype in addition to α 2 would increase antihyperalgesic efficacy and




Figure 1 | Distribution of the four subtypes of BDZ-sensitive GABA_ARs in the spinal cord. (a,b) Immunohistochemistry. (a) Localization of GABA_A α subunits in spinal cords of wt mice and of the four types of GABA_A triple point-mutated mice. Pseudocolour images display receptor density (lowest to highest density, blue to yellow). Right: quantitative comparison of expression levels in wt and triple point-mutated mice. Two-way analysis of variance (ANOVA) for interaction (α subunit \times genotype): $F(7,42) = 0.53$, $P = 0.68$, $n = 6$ mice (three sections each) for all groups, except RRHR, where $n = 7$ mice. Panel b same as a, but for $\gamma 2$ subunits. ANOVA followed by Dunnett's *post hoc* test. $F(4,15) = 0.32$; $P \geq 0.94$. $n = 4$ mice (three sections each) for all genotypes. (c,d) Autoradiography. (c) [^3H]Ro15-4513-binding sites indicate the distribution of all α/γ GABA_ARs independent of their BDZ sensitivity in wt mice and the four types of GABA_A triple point-mutated mice. H and R indicate homozygous wt and point-mutated alleles (for $\alpha 1/\alpha 2/\alpha 3/\alpha 5$ subunits), respectively. Right: quantification of [^3H]Ro15-4513-binding sites in the dorsal horn. ANOVA followed by Dunnett's *post hoc* test with wt as control. $F(4,10) = 2.89$, $P \geq 0.18$, $n = 3$, for all genotypes (>30 sections each). (d) Same as c, but distribution of BDZ-sensitive GABA_ARs labelled with [^3H]flumazenil. ANOVA followed by Dunnett's *post hoc* test with wt as control. $F(4,55) = 40.1$, $***P < 0.001$, $n = 11, 12, 12, 12$ and 14 mice (>10 sections each), for wt (HHHH), HRRR, RHRR, RRHR and RRRH mice, respectively. (e) Same as (d), but wt and GABA_A single point-mutated mice. $F(4,42) = 0.441$, $P > 0.66$, $n = 11, 9, 9, 9$ and 9 mice, for wt (HHHH), RHHH, HRHH, HHRH and HHHR mice, respectively. All scale bars, $200 \mu\text{m}$. All quantitative data mean \pm s.e.m.

compared antihyperalgesic responses in triple point-mutated mice with those in mice carrying only one or two point-mutated receptors. We found only small and statistically insignificant differences indicating that adding activity at a subtype different from $\alpha 2$ did not significantly increase antihyperalgesia (Fig. 2f).

The autoradiography data shown in Fig. 1d suggest a high prevalence of GABA_ARs with two different α subunits ('mixed GABA_ARs') in the spinal cord. Because the sum of the four receptor subtypes detected in the triple point-mutated mice greatly exceeded the total number of [^3H]flumazenil binding sites

in the wt mice, the pharmacological effect of a given GABA_A subtype may be overestimated when tested in triple point-mutated mice. To address this issue, we compared the level of antihyperalgesia achieved in the triple point-mutated mice with the loss in antihyperalgesia that occurs after mutation of the same subunit (Fig. 2g). For both types of sensory tests (Hargreaves and von Frey filament) and for all three α subunits under study, we found that the level of antihyperalgesia achieved in the triple point-mutated mice always exceeded the loss in antihyperalgesia by mutation of the same subunit consistent with the high

Table 1 | Baseline nociceptive sensitivities of all GABA_AR genotypes under study.

Genotype	BDZ-sensitive α subunit (-s)	Pre-CCI		Post-CCI	
		von Frey (g) (number of mice)	Hargreaves (s) (number of mice)	von Frey (g) (number of mice)	Hargreaves (s) (number of mice)
HHHH (wt)	$\alpha 1, \alpha 2, \alpha 3$ and $\alpha 5$	4.04 \pm 0.07 (20)	22.9 \pm 0.72 (17)	1.48 \pm 0.23 (8)	11.7 \pm 1.03 (16)
RHHH	$\alpha 2, \alpha 3$ and $\alpha 5$	4.41 \pm 0.11 (23)	22.6 \pm 0.63 (15)	1.52 \pm 0.11 (15)	9.68 \pm 0.81 (12)
RHHR	$\alpha 2$ and $\alpha 3$	4.21 \pm 0.12 (26)	23.4 \pm 0.57 (15)	1.46 \pm 0.09 (20)	10.4 \pm 1.15 (10)
RHRH	$\alpha 2$ and $\alpha 5$	4.05 \pm 0.09 (23)	22.9 \pm 0.60 (20)	1.26 \pm 0.10 (9)	12.8 \pm 1.70 (7)
RRHH	$\alpha 3$ and $\alpha 5$	4.23 \pm 0.04 (29)	21.0 \pm 0.43 (29)	1.52 \pm 0.06 (8)	8.54 \pm 0.62 (19)
RHRR	$\alpha 2$	4.12 \pm 0.06 (32)	22.2 \pm 0.53 (32)	1.42 \pm 0.11 (18)	10.7 \pm 0.72 (19)
RRHR	$\alpha 3$	4.12 \pm 0.12 (29)	22.2 \pm 0.50 (29)	1.27 \pm 0.08 (13)	10.3 \pm 1.00 (9)
RRRH	$\alpha 5$	4.32 \pm 0.07 (23)	21.5 \pm 0.42 (23)	1.67 \pm 0.08 (11)	10.2 \pm 0.89 (12)
RRRR	None	3.99 \pm 0.10 (9)	20.8 \pm 1.48 (6)	1.18 \pm 0.16 (9)	10.3 \pm 1.51 (6)
Statistics		F(8,205) = 1.57; $P \geq 0.06$	F(8,177) = 1.93; $P \geq 0.06$	F(8,102) = 1.78; $P \geq 0.16$	F(8,101) = 1.80; $P \geq 0.09$

ANOVA, analysis of variance; BDZ, benzodiazepine; CCI, chronic constriction injury; GABA_AR, GABA_A receptor. Values are given as mean \pm s.e.m. ANOVA followed by Dunnett's *post hoc* test with wt (HHHH) as control.

prevalence of mixed GABA_AR_s in the spinal cord. It is important to note that the rank order of antihyperalgesic efficacies was the same in experiments with triple point-mutated mice and in loss-of-function experiments in single and double point-mutated mice, and also identical to those found in a pain model employing chemical nociceptor activation instead of neuropathy (Fig. 2h).

Taken together, these results support a major contribution of $\alpha 2$ GABA_AR_s to antihyperalgesia in different pain models. However, because all experiments had to be conducted on an $\alpha 1$ point-mutated background to avoid confounding sedation, the relevance of $\alpha 2$ GABA_AR_s might be overestimated if ‘pain-relevant’ $\alpha 2$ GABA_AR_s contained also $\alpha 1$ subunits. To address this potential caveat we used HZ-166 (ref. 18), a novel benzodiazepine site ligand with an improved $\alpha 2/\alpha 1$ selectivity ratio¹⁹ and significant analgesic activity already at non-sedative doses²⁰ (Fig. 3). Antihyperalgesia by HZ-166 was almost completely blocked in single $\alpha 2$ (H101R) point-mutated (HRHH) mice, further confirming that the ‘pain-relevant’ GABA_AR_s exhibit an $\alpha 2$ pharmacology.

Potential undesired BDZ effects in triple point-mutated mice.

We next used the same triple point-mutated mouse approach to investigate non-pain-related effects. In these experiments, we focused on diminished locomotor activity (as a surrogate parameter of sedation), muscle strength and motor coordination. No significant differences were observed in the behaviour of drug-naïve wt and point-mutated mice (Table 2). The DZP treatment strongly reduced locomotor activity in wt mice and in mice with BDZ-sensitive GABA_AR_s of only the $\alpha 1$ subtype (HRRR mice; Fig. 4a). No sedative effects were observed in any of the other triple point-mutated mice. Mice with only $\alpha 2$ BDZ-sensitive GABA_AR_s (RHRR mice) even showed a strong increase in locomotor activity, which may originate from the anxiolytic effect of DZP occurring through $\alpha 2$ GABA_AR_s (ref. 12). Impairment of muscle strength was assessed in the horizontal wire test. Significant muscle relaxation was detected in wt mice and in mice with BDZ-sensitive GABA_AR_s of either only the $\alpha 2$ (RHRR mice) or $\alpha 3$ subtype (RRHR mice; Fig. 4b). Motor coordination, tested in the rotarod test, was significantly impaired by DZP in mice with only $\alpha 1$ (HRRR mice) and with only $\alpha 3$ BDZ-sensitive GABA_AR_s (RRHR mice; Fig. 4c). Quadruple point-mutated (RRRR) mice were completely protected from DZP-induced muscle relaxation and motor impairment. They did, however, show a trend towards reduced locomotor activity (compare Fig. 4a). We therefore assessed changes in locomotion also after higher DZP doses in the quadruple point-mutated mice and

found significant and dose-dependent impairment starting at 30 mg kg⁻¹ (Fig. 4d). Unlike $\alpha 1$ GABA_AR-mediated sedation, impairment of muscle strength was absent in the quadruple point-mutated mice even at doses ≥ 30 mg kg⁻¹. The sedative action of DZP remaining in the quadruple point-mutated mice may be attributed to a low-affinity BDZ-binding site at $\alpha 1$ GABA_AR_s described earlier^{21,22}.

Our findings on the contribution of $\alpha 1, \alpha 2$ and $\alpha 3$ GABA_AR_s to sedation, anxiolysis and muscle relaxation confirm previous studies using single GABA_AR point-mutated mice^{11,12,23}. The results from our experiments with triple point-mutated mice on motor coordination, however, differ from those obtained with single point-mutated mice^{11,12}. This discrepancy may arise from the involvement of mixed GABA_AR_s containing $\alpha 1$ and $\alpha 3$ subunits. We therefore tested whether motor coordination would also be impaired by TP003, an $\alpha 3$ GABA_AR-selective BDZ site agonist²⁴. Two hours after treatment with TP003 (10 mg kg⁻¹, p.o.), the time to fall off the rod was similarly decreased in wt and RRHR mice (by 43.9 \pm 10.2%, paired *t*-test $P < 0.05$, $n = 5$, in wt mice, and by 38.5 \pm 6.4, $n = 5$, $P < 0.05$), while HHRH and vehicle-treated wt mice showed no impairment (0.7 \pm 12.0%, $n = 5$, $P = 0.91$ and -4.7 \pm 9.2%, $n = 5$, $P = 0.59$).

Liability to tolerance development. Another major limitation of classical BDZs is their liability to tolerance development (that is, the loss of activity during prolonged use). We tested whether this tolerance would also occur for the antihyperalgesic actions of BDZs and whether it could be prevented by selectively targeting $\alpha 2$ GABA_AR_s. To this end, we applied again CCI surgery and treated mice of the different strains for nine consecutive days with 10 mg kg⁻¹ DZP or vehicle p.o. once daily. On the 10th day, both groups of mice were split into two subgroups, treated with either 10 mg kg⁻¹ DZP or vehicle, and mechanical hypersensitivity was monitored for 2 h after drug application. We first did this experiment in single $\alpha 1$ point-mutated (RHHH) mice to avoid confounding sedation (Fig. 5a,d). While DZP-naïve mice responded with an almost complete reversal of hyperalgesia, mice chronically pretreated with DZP showed no antihyperalgesic response. We then performed this experiment in triple point-mutated RHRR mice with only $\alpha 2$ BDZ-sensitive GABA_AR_s and found that these mice were completely protected from tolerance development (Fig. 5b,d). In the latter experiment, the antihyperalgesic response was generally lower than in $\alpha 1$ point-mutated (RHHH) mice. We therefore repeated the first experiment with a lower dose of DZP (3 mg kg⁻¹) to exclude the possibility that the absence of tolerance in the RHRR mice was




Figure 2 | Antihyperalgesia by per oral DZP in GABA_AR point-mutated mice. (a) Reversal of mechanical hyperalgesia by DZP in $\alpha 1$ point-mutated (RHHH) mice assessed with von Frey filaments 7 days after CCI surgery. (b) Dose-response relationship. Data were fitted to Hill's equation. *** $P < 0.001$; ** $P < 0.01$ versus vehicle (ANOVA $F(4,28) = 29.0$ followed by Dunnett's *post hoc* test, $n = 7, 7, 6, 7$ and 6 mice, for vehicle and $1, 10$ and 30 mg kg^{-1} DZP, respectively). (c,d) Reversal of mechanical (c) and heat hyperalgesia (d) by DZP (10 mg kg^{-1}) 1 week after the CCI surgery. All mice except wt carried the H→R point mutation in $\alpha 1$ GABA_ARs to avoid sedation. (e) Comparison of antihyperalgesia (determined from 70 to 130 min after DZP administration) in GABA_AR triple point-mutated mice. *** $P < 0.001$; ** $P < 0.01$; versus vehicle-treated wt mice (ANOVA followed by Dunnett's *post hoc* test, $F(4,54) = 36.2$ and $F(4,53) = 15.1$ for von Frey and Hargreaves tests, respectively). No antihyperalgesia by DZP occurred in quadruple GABA_AR point-mutated mice ($P = 0.85$ and 0.94). (f) Comparison of antihyperalgesia by DZP in RHRR mice with that of mice carrying, in addition, BDZ-sensitive $\alpha 3$ and/or $\alpha 5$ GABA_ARs. No significant differences were observed (ANOVA followed by Dunnett's *post hoc* test, von Frey: $F(3,57) = 2.96$; $P \geq 0.12$. Hargreaves: $F(3,44) = 0.98$; $P \geq 0.37$). (g) Antihyperalgesia in triple GABA_AR point-mutated mice with only one DZP-sensitive GABA_AR subtype (grey columns) compared with antihyperalgesia lost through H→R point mutation of the same subunit (light red columns). Antihyperalgesia in triple point-mutated mice is expressed as percentage of antihyperalgesia in RHHH mice. Loss of antihyperalgesia was calculated as the reduction in antihyperalgesia relative to RHHH mice. *** $P < 0.001$; ** $P < 0.01$; * $P < 0.05$ (unpaired *t*-tests). Number of mice in c–g: vehicle-treated wt (von Frey/Hargreaves test) $n = 7/11$, RHHH $n = 15/12$, RHRR $n = 18/19$, RRHR $n = 13/9$, RRRH $n = 11/12$, RHHH $n = 19/10$, RHRR $n = 9/7$, RRRH $n = 8/19$. (h) Formalin test. Reduction by DZP (10 mg kg^{-1}) in the number of flinches and licking bouts compared with vehicle-treated mice. *** $P < 0.001$; ** $P < 0.01$; * $P < 0.05$ (ANOVA followed by Dunnett's *post hoc* test. Flinches, $F(3,24) = 6.71$; licking bouts, $F(4,25) = 10.7$. $n = 9, 6, 7$ and 6 mice, for vehicle-treated wt and DZP-treated RHRR, RRHR and RRRH mice, respectively. All data points are mean \pm s.e.m.

due to a smaller DZP effect. With this lower dose, complete tolerance still developed in RHHH mice (Fig. 5c,d). These experiments indicate that activity at $\alpha 3$ or $\alpha 5$ GABA_ARs was necessary to induce antihyperalgesic tolerance. We therefore tested double point-mutated mice in which $\alpha 3$ GABA_ARs or

$\alpha 5$ GABA_ARs were left BDZ-sensitive in addition to $\alpha 2$ GABA_ARs (RHHR and RHRH mice). Complete tolerance developed in RHHR mice, but not in RHRH mice, indicating that activity at $\alpha 3$ GABA_ARs was necessary for induction of antihyperalgesic tolerance in mutated mice (Fig. 5e). In this set of experiments, we




Figure 3 | Antihyperalgesia by the non-sedative BDZ HZ-166 occurs through $\alpha 2$ GABA_ARs. (a) HZ-166 (16 mg kg⁻¹, i.p.) almost completely reversed CCI-induced mechanical sensitization in wt mice (HHHH, black), but not in $\alpha 2$ (H101R) mice (HRHH mice, red). (b) Statistical analysis. ***P* < 0.01 (unpaired *t*-test), *n* = 7 and 8 mice, for wt and $\alpha 2$ (H101R) mice. All data points are mean \pm s.e.m.

finally tested whether the same subtype dependence would also be found for tolerance against $\alpha 2$ GABA_AR-mediated anxiolysis (Fig. 5f). Unlike antihyperalgesic tolerance, anxiolytic tolerance (measured as increased locomotor activity in the open field) still developed in RHRR mice. This dissociation suggests that tolerance development against antihyperalgesia is not a receptor or cell-autonomous process²⁵ but rather occurs at the circuit level.

Why classical BDZs lack clinically relevant analgesic properties. Our present results and previous pain studies employing subtype-selective BDZ site agonists in rodents^{5,20,26–29} contrast with the lack of a clear analgesic or antihyperalgesic action of classical BDZs in human patients^{30,31}. Apart from species differences and differences between disease models and actual disease in human patients, we found one possible explanation particularly worth studying. The doses and the degrees of receptor activation required for a relevant effect might be significantly higher in case of antihyperalgesia than of sedation. As a consequence, antihyperalgesia would occur in patients only at doses already inducing strong sedation. The availability of the triple point-mutated mice allowed us to directly compare the

Table 2 Baseline locomotor activity and performance in the horizontal wire and rotarod tests.				
Genotype	BDZ-sensitive α subunit (-s)	Activity counts (number of mice)	Horizontal wire performance* (number of mice)	Rotarod performance† (number of mice)
HHHH (wt)	$\alpha 1$, $\alpha 2$, $\alpha 3$ and $\alpha 5$	962 \pm 114 (14)	92 \pm 2.5 (17)	92 \pm 5.6 (5)
HRRR	$\alpha 1$	1102 \pm 104 (9)	98 \pm 1.2 (9)	98 \pm 8.2 (6)
RHRR	$\alpha 2$	1132 \pm 106 (8)	96 \pm 1.2 (8)	81 \pm 5.3 (5)
RRHR	$\alpha 3$	1279 \pm 114 (10)	99 \pm 1.1 (10)	85 \pm 7.7 (5)
RRRH	$\alpha 5$	987 \pm 90.4 (8)	100 \pm 0.0 (8)	84 \pm 11 (5)
RRRR	None	1037 \pm 115 (9)	99 \pm 1.2 (9)	110 \pm 13 (5)
Statistics‡		F(5,52) = 1.16; <i>P</i> \geq 0.13	F(5,55) = 2.10; <i>P</i> \geq 0.70	F(5,25) = 1.53; <i>P</i> \geq 0.47

ANOVA, analysis of variance; BDZ, benzodiazepine; wt, wild type.
Values are given as mean \pm s.e.m.
*Horizontal wire performance, expressed as success rate (%).
†Rotarod performance, expressed as time before fall off (s).
‡ANOVA followed by Dunnett's *post hoc* test with wt (HHHH) as control.




Figure 4 | Sedation, muscle relaxation and motor coordination in GABA_AR-mutated mice. Effects of DZP (10 mg kg⁻¹, p.o.) on locomotor activity in the open field test (a), on muscle relaxation in the horizontal wire test (b), and on motor coordination in the rotarod test (c). ****P* < 0.001, **P* < 0.05 significant versus vehicle-treated wt (HHHH) mice (ANOVA followed by Dunnett's *post hoc* test). Statistics: locomotor activity F(6,177) = 79.5 (*n* = 106, 26, 8, 8, 9, 13 and 13 mice, for vehicle and DZP-treated wt mice, and DZP-treated HRRR, RHRR, RRHR, RRRH and RRRR mice, respectively). Horizontal wire F(6,165) = 44.0 (*n* = 109, 14, 8, 8, 9, 13 and 10 mice). Rotarod F(6,41) = 11.5 (*n* = 8, 5, 6, 7, 6, 8 and 8 mice). (d) Effects of DZP on locomotor activity and horizontal wire performance in quadruple GABA_AR point-mutated (RRRR) mice. ****P* < 0.001; ***P* < 0.05 significant versus vehicle (ANOVA followed by Dunnett's *post hoc* test) F(3,33) = 13.4 (locomotor activity), *n* = 9, 13, 8 and 7 mice, for vehicle, and 10, 30 and 100 mg kg⁻¹ DZP; F(3,30) = 0.44; *P* > 0.60 (horizontal wire test), *n* = 9, 10, 8 and 7 mice, for vehicle, and 10, 30 and 100 mg kg⁻¹ DZP. All data points are mean \pm s.e.m.




Figure 5 | Tolerance liability against the antihyperalgesic effects of DZP. Starting on day 5 after the CCI surgery, mice were treated with DZP or vehicle once daily for nine consecutive days. On day 10, mice were given either DZP or vehicle, and antihyperalgesic effects were measured for 2 h. **(a)** Complete loss of antihyperalgesic activity was observed after 9-day DZP treatment (10 mg kg⁻¹, p.o.) in RHHH mice. **(b)** Same as **a**, but RHRR triple point-mutated mice. These mice were completely protected from tolerance development against DZP-induced antihyperalgesia. **(c)** Same as **a**, but lower dose of DZP (3 mg kg⁻¹, p.o.). Tolerance in RHHH mice still developed at a lower dose of DZP, which induced an antihyperalgesic effect similar to that of 10 mg kg⁻¹ in RHRR mice. **(d)** Statistical comparison. Two-way ANOVA for the interaction pretreatment \times acute treatment ($F(3,25) = 32.5$ ($n = 5, 8, 7$ and 8 mice for veh/veh, veh/DZP, DZP/veh and DZP/DZP groups, respectively); $F(3,25) = 0.014$ ($n = 7, 8, 7$ and 6); $F(3,24) = 5.97$ ($n = 8, 7, 7$ and 6), for data shown in **a–c**). Maximum antihyperalgesic activity was calculated for the interval between 80 and 120 min after DZP administration. **(e)** Double GABA_AR point-mutated mice (RHHR and RHRH), in which $\alpha 3$ GABA_AR or $\alpha 5$ GABA_AR were left BDZ-sensitive in addition to $\alpha 2$ GABA_AR. Tolerance development required the additional presence of DZP-sensitive $\alpha 3$ GABA_AR. Two-way ANOVA for the interaction pretreatment \times acute treatment. RHHR mice: $F(3,22) = 22.1$ ($n = 6, 6, 7$ and 6 mice for veh/veh, veh/DZP, DZP/veh and DZP/DZP groups, respectively); RHRH mice: $F(3,21) = 0$ ($n = 6$ for all groups). **(f)** Unlike antihyperalgesic effects, anxiolytic effects of DZP were still susceptible to tolerance development in RHRR mice after 9-day treatment with DZP (10 mg kg⁻¹, p.o.). Two-way ANOVA $F(3,26) = 28.3$ ($n = 8, 7, 7$ and 7) for the interaction pretreatment \times acute treatment. *** $P = 0.001$; * $P = 0.05$. All data points are mean \pm s.e.m.

doses and levels of receptor occupancy at the relevant GABA_AR subtypes and sites. For these experiments, we chose midazolam (MDZ) as a second classical BDZ in addition to DZP. First, we determined for both drugs their $\alpha 2/\alpha 1$ selectivity profiles in electrophysiological experiments on recombinant GABA_ARs. As expected, we found that DZP potentiated $\alpha 1/\beta 2/\gamma 2$ and $\alpha 2/\beta 3/\gamma 2$ GABA_ARs with similar efficacy and potency (Fig. 6a and Table 3). By contrast, MDZ potentiated $\alpha 1/\beta 2/\gamma 2$ GABA_ARs more than twice as much as $\alpha 2/\beta 3/\gamma 2$ GABA_ARs (Fig. 6b and Table 3). We then verified that the H \rightarrow R point mutation blocked not only DZP effects but also MDZ binding and GABA_AR potentiation (Figs 6c,d; see also ref. 32). Next, we compared the dose dependency of DZP- and MDZ-induced antihyperalgesia in mice with only $\alpha 2$ BDZ-sensitive GABA_ARs (RHRR mice) with that of

DZP- and MDZ-induced sedation in mice with only $\alpha 1$ BDZ-sensitive GABA_ARs (HRRR mice). We found that half maximal sedation occurred already at a doses of 0.33 ± 0.05 and 0.52 ± 0.11 mg kg⁻¹ (mean \pm s.d.) for DZP and MDZ, respectively, while half maximal antihyperalgesia required 8 ± 6 mg kg⁻¹ (DZP) and 10.4 ± 2.0 mg kg⁻¹ (MDZ). A rightward shift of the response curve was also observed when the degrees of receptor occupancy required for antihyperalgesia and for sedation were compared. Half maximal sedation was reached when DZP had bound $47 \pm 6\%$ brain $\alpha 1$ GABA_ARs, while half maximal antihyperalgesia required $71 \pm 2\%$ occupancy at spinal $\alpha 2$ GABA_ARs. In case of MDZ, the required receptor occupancies were even further apart ($24 \pm 2\%$ of brain $\alpha 1$ GABA_ARs and $71 \pm 2\%$ of spinal $\alpha 2$ GABA_ARs).




Figure 6 | Activity of DZP and MDZ at $\alpha 1$ and $\alpha 2$ GABA_ARs, and PK/PD modelling. (a,b) Potentiation of recombinant $\alpha 1\beta 2\gamma 2$ and $\alpha 2\beta 3\gamma 2$ GABA_A currents by 1 μ M DZP (a) and 1 μ M MDZ (b). GABA concentration was EC₅ (5 μ M for $\alpha 1\beta 2\gamma 2$ and 2 μ M for $\alpha 2\beta 3\gamma 2$). $n = 6$ for both drugs and all concentrations. Scales bars, 200 pA, 2 s. (c,d) The H101R mutation prevented potentiation of $\alpha 1\beta 2\gamma 2$ GABA_A currents by MDZ (** $P < 0.01$, $n = 5$, unpaired t -test; c) and binding of MDZ to $\alpha 1\beta 2\gamma 2$ GABA_ARs ($n = 6$ for all concentrations in wt and mutated receptors; d). (e-h) $\alpha 1$ GABA_A-mediated sedation and $\alpha 2$ GABA_A-mediated antihyperalgesia by DZP and MDZ. Sedation (black symbols) in HRRR mice was assessed in the open field test and expressed as percent maximum possible effect determined from the reduction in activity compared with vehicle-treated mice of the same genotype. Antihyperalgesia (red symbols) in RHRR mice was determined as the increase in mechanical withdrawal thresholds compared with pre-drug values 1 week after the CCI surgery. (e,f) Dose-response relationships. Data were fitted to Hill's equation with baseline fixed at 0. Sedation and antihyperalgesia were determined between 15 and 90 min after oral DZP (e) or between 15 and 60 min after oral MDZ (f). Mice were killed immediately afterwards (that is, at the time point of maximal effects), and brains and spinal cords were removed for further analyses. (g,h) Dependence on receptor occupancy (RO). (g) Sedation versus RO in brains of HRRR mice treated with DZP (0.03, 0.1, 0.3, 1, 3, 10 and 30 mg kg^{-1} , $n = 6$ mice for all doses). Antihyperalgesia versus RO in lumbar spinal cords of RHRR mice treated with 0.1, 1, 3, 10 or 30 mg kg^{-1} DZP, $n = 6$ mice for all doses. (h) Same as g but MDZ. Sedation 0.1 ($n = 6$ mice), 0.3 (7), 0.5 (7), 1.0 (7), 3 (6) and 10 (4) mg kg^{-1} . Antihyperalgesia 0.1 ($n = 3$), 0.3 (5), 1.0 (9), 3 (8), 5 (7) and 10 (7) mg kg^{-1} MDZ. Data were fitted to sigmoidal functions. Data shown in e,g (on DZP) and in f,h (on MDZ) are from the same two groups of mice. All data points are mean \pm s.e.m.

consistent with the even less favourable $\alpha 2/\alpha 1$ selectivity ratio of MDZ. These data indicate that, when applied to wt mice, the DZP and MDZ doses needed for half maximal antihyperalgesia induce an almost complete ($\sim 95\%$) reduction in locomotor activity, while at non-sedative doses both BDZs would not induce significant antihyperalgesia. Dose-limiting sedation is therefore the most likely reason for the absence of a relevant antihyperalgesic activity of classical, nonselective BDZs in human patients.

Discussion

In the present study we have employed triple GABA_A point-mutated mice to characterize the pharmacological

actions expected from yet-to-be-developed subtype-selective BDZ-binding site agonists. Our study can be viewed as a 'restriction-of-function' approach ('what effect of DZP remains when only a single GABA_A subtype is targeted?') as opposed to previous loss-of-function studies in single GABA_A point-mutated mice ('which effects of DZP are lost or reduced when activity at one GABA_A subtype is abolished compared with DZP-treated wt mice?'). With respect to sedation, anxiolysis and the muscle-relaxant action, the present study confirms previous results obtained with single point-mutated mice: sedative actions of BDZs occur through $\alpha 1$ GABA_A (refs 10,11), anxiolytic effects through $\alpha 2$ GABA_ARs (ref. 12) and muscle relaxation through $\alpha 2$ and $\alpha 3$ GABA_A (ref. 23).

Table 3 | PK/PD parameters* of DZP and MDZ.

	DZP		MDZ	
	$\alpha 1/\beta 2/\gamma 2$	$\alpha 2/\beta 3/\gamma 2$	$\alpha 1/\beta 2/\gamma 2$	$\alpha 2/\beta 3/\gamma 2$
EC ₅₀ (μM)	0.081 ± 0.02	0.15 ± 0.04	0.17 ± 0.18	0.092 ± 0.019
E _{max} (% pot I _{GABA})	91.6 ± 7.7	103 ± 11	235 ± 43	75.8 ± 4.7
n _H	1.50 ± 0.60	1.20 ± 0.35	0.73 ± 0.26	1.23 ± 0.29
α2/α1 selectivity†	1.25		0.32	
ED ₅₀ (mg kg ⁻¹)	Sedation 0.33 ± 0.05	Analgesia 7.6 ± 0.6	Sedation 0.52 ± 0.11	Analgesia 10.4 ± 2.0
E _{max} (%MPE)	93.6 ± 4.5	100 (Fixed)	92.2 ± 10.4	100 (Fixed)
n _H	2.0 ± 0.64	0.75 ± 0.19	2.28 ± 0.98	1.42 ± 0.40
ED ₅₀ (sedation/analgesia)	0.043		0.050	
RO ₅₀ (%)	Sedation 46.9 ± 5.5	Analgesia 70.9 ± 1.7	Sedation 23.7 ± 2.2	Analgesia 76.5 ± 1.8
E _{max}	97.9 ± 7.1	100 (Fixed)	94.7 ± 11.8	100 (Fixed)
Rate	14.1 ± 4.1	14.9 ± 2.0	4.14 ± 2.10	5.01 ± 2.11
RO ₅₀ (sedation/analgesia)	0.66		0.31	

PK, pharmacokinetic; PD, pharmacodynamic; DZP, diazepam; MDZ, midazolam; %MPE, percent maximal possible effect; RO, receptor occupancy.
For number of mice per group see Fig. 6.
*Values are mean ± s.d.
†E_{max} (α2/β3/γ2)/E_{max} (α1/β2/γ2).

Discrepancies between results from restriction-of-function and loss-of-function approaches were observed for impairment of motor coordination. The present study shows that undesired impairment of motor coordination is caused by DZP in HRRR and RRHR mice (that is, it is evoked when either α1 or α3GABA_ARs are specifically targeted), while previous reports in the respective single point-mutated mice failed to provide evidence for an involvement of these GABA_AR subtypes^{11,12}. This discrepancy is consistent with the idea that DZP impairs motor coordination through mixed GABA_ARs containing an α1 and an α3 subunit (α1/α3GABA_ARs). Immunohistochemical analyses have shown that co-expression of α1 and α3 GABA_AR subunits occurs in subsets of central neurons^{33,34}, and biochemical data have provided direct evidence for the presence of different α subunits within the same GABA_AR complex¹⁵. In particular, α3 containing GABA_ARs occur mainly as mixed α1/α3GABA_ARs (refs 14,15). Furthermore, biochemical data also suggest that in mixed GABA_ARs with one point-mutated α subunit the nonpoint-mutated (wt) subunit has a higher probability for interaction with the γ2 subunit than the point-mutated α subunit^{15,16}. Results from loss-of-function analysis in single point-mutated mice and from restriction-of-function analyses in triple point-mutated mice should therefore be seen as lower and upper limits for the contribution of a given GABA_AR subtype to an *in vivo* drug action. Conversely, the lack of sedation in α1 point-mutated mice^{10,11} and of anxiolysis in α2 point-mutated mice¹² suggests that these actions occur through GABA_ARs with only a single type of α subunit. In line with this conclusion is that the majority of GABA_ARs in the brain are pure α1GABA_ARs (α1/α1GABA_ARs)¹⁵. Similarly, BDZ-mediated anxiolysis occurs through pure α2GABA_ARs, with α2/α2GABA_ARs being the single most prevalent isoform of α2 containing GABA_ARs at supraspinal sites¹⁵.

As demonstrated by our autoradiography experiments (compare Fig. 1d,e), mixed GABA_ARs make up the vast majority of GABA_ARs in the spinal cord. Following the model described above, one would expect that in case of mixed GABA_ARs the single point mutation approach would yield smaller effects than the triple point mutation strategy. For instance, in case of an αx/αyGABA_AR neither the αx single point mutation nor the αy single point mutation would abolish BDZ sensitivity; however, both triple point-mutated mice with only αx or only αy

remaining BDZ-sensitive would yield full behavioural effects. Results for antihyperalgesia obtained in the present study were consistent with a contribution of mixed GABA_ARs to antihyperalgesia. For all three contributing α subunits, the loss-of-function obtained through point mutation of one α subunit was smaller than the restriction-of-function in the respective triple point-mutated mouse. The discrepancies were particularly large for α3 in mechanical sensitization and for α5 in heat hyperalgesia, suggesting that these effects occur mainly through mixed GABA_ARs. Importantly, the rank orders of antihyperalgesic efficacy were the same for the restriction-of-function and loss-of-function approaches with α2 > α5 > α3 for mechanical sensitization and α2 > α3 > α5 for heat hyperalgesia and chemical nociception. The same rank order of efficacies has also been reported previously for antihyperalgesia using local spinal injections in single GABA_AR point-mutated mice⁵. Our present results therefore corroborate the critical importance of α2GABA_ARs as targets for antihyperalgesia.

Our experiments on the quadruple point-mutated mice indicate that antihyperalgesia, anxiolysis, muscle relaxation and impairment of motor coordination occur through the high-affinity BDZ-binding site formed by the α/γ interface. Among the DZP actions assessed here, only sedation by high doses of DZP occurred in quadruple point-mutated mice. This result is consistent with previous electrophysiological data, which suggested the presence of a low-affinity BDZ-binding site in α1GABA_ARs contributing to the anaesthetic actions of BDZs²².

Most BDZ effects quickly diminish during prolonged treatment (that is, they undergo fast tolerance development). In the present study, we show that this is also the case for antihyperalgesia. Previous studies with the non-sedative BDZ site ligands L-838,417 and HZ-166 showed that these compounds have strongly reduced liabilities to tolerance development^{5,20,27}. It has, however, not been possible to attribute this reduced tolerance to generally reduced agonistic activity or to improved subtype specificity. Our present study shows that tolerance can be avoided when only α2GABA_ARs are targeted, even with compounds that exert full agonistic activity. Additional experiments in RHRH and RRHR mice exclude α5GABA_ARs and suggest that activity at α3GABA_ARs is required for tolerance induction. However, because all these experiments were carried out in mice carrying the H → R point mutation in the α1 subunit, we cannot exclude

that tolerance occurs through mixed $\alpha 1/\alpha 3$ GABA_ARs, which in the wt situation may exhibit an $\alpha 1$ pharmacology. Such a scenario would explain why compounds with activity at $\alpha 3$ GABA_ARs but absent or reduced activity at $\alpha 1$ GABA_ARs lack liability to tolerance^{5,20,27}. Interestingly, tolerance against the anxiolytic action of DZP was retained in triple point-mutated RHRR mice with only $\alpha 2$ GABA_ARs remaining BDZ-sensitive. This difference suggests that tolerance development against the different BDZ effects involves activity at distinct subunits and possibly different mechanisms. Cell- or receptor-autonomous processes, which have been proposed for the desensitization of hippocampal $\alpha 2$ GABA_ARs (ref. 25), are unlikely to be responsible for the tolerance against antihyperalgesia.

Addictive (reinforcing) properties of classical BDZs are another area of concern that has not been addressed in the present study. Reduced BDZ-induced reward facilitation has been reported in mice carrying the H→R point mutation in either the $\alpha 1$, $\alpha 2$ or $\alpha 3$ GABA_AR subunits^{35,36}. Preference for MDZ in a two-bottle choice paradigm was absent in $\alpha 1$ (refs 31,35) and $\alpha 2$ point-mutated mice³⁵, suggesting that the simultaneous positive modulations of both the $\alpha 1$ and $\alpha 2$ GABA_AR are required for reinforcement. Tan *et al.*³² reported that $\alpha 1$ GABA_ARs on GABAergic neurons in the ventral tegmental area (VTA) are required for MDZ-induced neuronal plasticity, strengthening glutamatergic excitation in the VTA. These findings suggest that activity at more than one GABA_AR subtype is necessary for BDZ reinforcement. $\alpha 2$ -selective BDZ site agonists might therefore be largely devoid of addictive properties.

Previous studies have provided evidence that activation of supraspinal GABA_ARs might enhance pain and counteract spinal antihyperalgesia, for example, through the inhibition of descending antinociceptive fibre tracts^{37–39}. In our study, we did not observe any pronociceptive actions of systemically administered DZP. If such pronociceptive actions were relevant in the pain models used here, they would occur through $\alpha 1$ GABA_ARs whose functions in nociception were not addressed in the present study. Any such effects would be avoided by $\alpha 1$ -sparing BDZ site agonists.

The availability of the triple GABA_AR point-mutated mice permitted a direct pharmacokinetic/pharmacodynamic comparison of $\alpha 1$ -mediated sedation and $\alpha 2$ -mediated antihyperalgesia in the absence of confounding behavioural effects from other GABA_AR subtypes. In case of DZP, ED₅₀ values for sedation and antihyperalgesia differed by a factor of more than 20, and a 50% higher degree of receptor occupancy was needed for $\alpha 2$ -mediated antihyperalgesia compared with $\alpha 1$ -mediated sedation. New subtype-selective agents will therefore have to have a high degree of $\alpha 2$ over $\alpha 1$ selectivity in order to achieve clinically relevant antihyperalgesia in the absence of sedation. These data also indicate that dose-limiting sedation most likely underlies the lack of clinically relevant antihyperalgesic effects of presently used (nonselective) BDZs. This is consistent with a recent clinical study in human volunteers showing weak antihyperalgesic effects at doses that caused only mild sedation⁴⁰.

Which GABA_AR subtype should be targeted for an optimal benefit-risk ratio in pain treatment? Both the present restriction-of-function and loss-of-function experiments on systemic BDZ administration and previous experiments with local spinal injections and single GABA_AR point-mutated mice⁵ attribute the highest antihyperalgesic efficacy to $\alpha 2$ GABA_ARs. The present study has shown that adding activity at $\alpha 3$ GABA_ARs or $\alpha 5$ GABA_ARs increases antihyperalgesic efficacy only moderately or not at all. Because it is at present not known how the different mixed GABA_ARs respond to subtype-selective agents, a conservative prediction should be made on the basis of the results obtained from triple point-mutated mice. These experiments indicate that sedation is exclusively due to

activation $\alpha 1$ GABA_ARs, which is also consistent with studies using the subtype-selective agonist TPA023B that completely lacks agonistic activity at $\alpha 1$ GABA_ARs as well as sedation in humans⁴¹. The present experiments also showed that impaired motor coordination involves $\alpha 1$ GABA_ARs and/or $\alpha 3$ GABA_ARs. None of the undesired effects investigated in the present study could be attributed to $\alpha 5$ GABA_ARs. However, cognitive impairment by nonselective BDZ likely originates from activity at $\alpha 5$ GABA_ARs, as $\alpha 5$ GABA_AR-selective inverse agonists enhance cognitive capabilities⁴². Subtype-selective BDZ agonists targeting only $\alpha 2$ GABA_ARs should therefore have the best benefit-risk ratio. They will produce pronounced antihyperalgesia in the absence of sedation and will not interfere with motor coordination and should not lose antihyperalgesic activity during prolonged treatment. They will however have anxiolytic properties (at least during the beginning of the treatment) and muscle-relaxant effects. Both of these actions may be beneficial in chronic pain patients. Because the side effects studied here are relevant for other indications, we expect that our present findings will benefit not only the development of innovative subtype-selective BDZs as analgesics but also as drugs for the treatment of several prevalent psychiatric diseases⁴³.

Methods

Mice. Experiments were performed in wt mice, and in homozygous single, double, triple and quadruple (H→R) GABA_AR point-mutated mice, that is, in mice expressing different combinations of BDZ-sensitive and BDZ-insensitive GABA_AR α subunits. All mice were of the same genetic background (129X1/SvJ). Double, triple and quadruple point-mutated mice were generated by cross-breeding single point-mutated mice described earlier^{11,12,44}.

Authoradiography. The distribution of BDZ-sensitive GABA_AR subtypes in the lumbar spinal cord was examined in 16- μ m-thick horizontal sections, which were cut from fresh-frozen spinal cords. Sections were incubated with 5 nM [³H]flumazenil (50 Ci mmol⁻¹) or 9 nM [³H]Ro15-4513 (22.7 Ci mmol⁻¹) diluted in 50 mM Tris pH 7.4 for 120 min on ice. After washing three times for 20 s in ice-cold buffer, sections were dried and exposed along with [³H]-standards to a tritium-sensitive phosphorimaging screen (Cyclone Storage Phosphor Screen, Perkin Elmer). Quantification was carried out using the Optiquant software (Perkin Elmer). Nonspecific binding was assessed by co-incubating 10 μ M clonazepam ([³H]flumazenil binding) or 10 μ M flumazenil ([³H]Ro15-4513-binding).

Immunohistochemistry. The localization of GABA_AR subunits was visualized on 40- μ m-thick lumbar spinal cord cryosections by DAB immunoperoxidase staining on sections from artificial cerebrospinal fluid (aCSF)-perfused mice postfixed for 90 min in 4% paraformaldehyde (PFA) (without picric acid)⁴⁵. Antibodies were home-made subunit-specific antisera⁴⁵. Final dilutions were 1:20,000 ($\alpha 1$), 1:1,000 ($\alpha 2$), 1:10,000 ($\alpha 3$), 1:3,000 ($\alpha 5$) and 1:10,000 ($\gamma 2$). Images were taken with a bright field light microscope connected to a digital camera and processed with the AxioVision Rel. 4.5 software where an intensity-gradient false-colour filter was applied.

Behavioural experiments. All behavioural experiments were performed in 7- to 12-week-old female and male mice. Care was taken to ensure equal numbers of female and male mice. All behavioural experiments were made by a male experimenter, blinded either to the genotype of the mice or to their treatment with drug or vehicle. Permission for animal experiments was obtained from the Veterinäramt des Kantons Zürich (licence numbers 135/2009 and 126/2012).

DZP (suspended in 0.9% saline and 1% Tween80) and MDZ (dissolved in 0.9% saline pH 3.0) were applied orally in all experiments. HZ-166 (suspended in 0.5% methyl cellulose) was given intraperitoneally (i.p.).

Neuropathic pain was evoked through CCI¹⁷ of the left sciatic nerve proximal to the trifurcation with three loose (5–0 silk) ligatures. Mice, which showed signs of paralysis or which did not develop significant hypersensitivity, were excluded from subsequent experiments. Effects of DZP and MDZ (p.o. in 0.9% saline and 1% Tween 80) on thermal and mechanical hyperalgesia were assessed 1 week after surgery. Heat hyperalgesia was quantified as the change in the latency of paw withdrawal evoked by exposure of the plantar side of one hindpaw to a defined radiant heat stimulus. Mechanical hyperalgesia was assessed with electronic von Frey filament no. 7 (ITC, Woodland Hills, CA). Three to four measurements were made for each time point and animal for both heat and mechanical hyperalgesia. Percent maximal possible effect (%MPE) was calculated as follows:

$\%MPE(t) = (E(t) - E_{\text{predrug}}) / (E_{\text{preCCI}} - E_{\text{predrug}}); E(t)$, paw withdrawal thresholds or latency at time point t . E_{predrug} , E after CCI surgery but before DZP application; E_{preCCI} , E baseline before CCI surgery.

Formalin (4%, 20 μ l) was injected subcutaneously into the plantar surface of the left hindpaw. Flinches and licks of the injected paw were counted for 1 h in 5-min intervals starting immediately after formalin injection. DZP was administered p.o. 60 min before formalin.

Locomotor activity was measured in an open field arena (10 cm diameter) equipped with four pairs of light beams and photosensors. Mice were placed into the arena 75 min before DZP application. Locomotor activity was analysed for the time interval between 20 and 165 min after DZP administration.

Motor control was assessed with a rotarod accelerating from 4 to 40 r.p.m. within 5 min. Sixty minutes after DZP administration mice were placed on the rotarod. Five measurements were taken per mouse.

To assess muscle relaxation, mice were placed with their forepaws on a metal horizontal wire placed 20 cm above ground. Successes and failures to grab the wire with at least one hindpaw were recorded between 60 and 180 min after DZP administration.

Electrophysiology. The effects of DZP and MDZ on currents through recombinant GABA_ARs were studied in HEK293 cells (ATCC) transiently expressing GABA_ARs. HEK293 cells were transfected using lipofectamine LTX⁴⁶. To ensure expression of the $\gamma 2$ subunit (required for modulation of GABA_ARs by BDZs) in all recorded cells, we transfected cells with a plasmid expressing the $\gamma 2$ subunit plus enhanced green fluorescent protein from an internal ribosomal entry site (IRES), and only selected eGFP-positive cells for recordings. The transfection mixture contained (in μ g): 1 $\alpha 1$, 1 $\beta 2$ and 3 $\gamma 2$ /eGFP (used as a marker of successful transfection) or 1 $\alpha 2$, 1 $\beta 3$ and 3 $\gamma 2$ /eGFP. Recordings were made at 18–36 h after transfection. Whole-cell patch-clamp recordings of GABA-evoked currents were made at room temperature (20–24 °C) and at a holding potential of -60 mV. Recording electrodes were filled with solution containing (in mM): 120 CsCl, 10 EGTA, 10 HEPES (pH 7.40), 4 MgCl₂, 0.5 GTP and 2 ATP. The external solution contained (in mM): 150 NaCl, 10 KCl, 2.0 CaCl₂, 1.0 MgCl₂, 10 HEPES (pH 7.4) and 10 glucose. GABA was applied to the recorded cell using a manually controlled pulse (4–6 s) of a low subsaturating GABA concentration (EC_{50}). EC_{50} values of GABA were determined for wt $\alpha 1\beta 2\gamma 2$, wt $\alpha 2\beta 3\gamma 2$ and mutant $\alpha 1(H101R)\beta 2\gamma 2$ receptors. EC_{50} values and Hill coefficients (n_H) were obtained from fits of normalized concentration–response curves to the equation $I_{\text{GABA}} = I_{\text{max}} [GABA]^{n_H} / ([GABA]^{n_H} + [EC_{50}]^{n_H})$. I_{max} was determined as the average maximal current elicited by a concentration of 1 mM GABA. DZP and MDZ dissolved in DMSO and subsequently diluted with recording solution were co-applied together with GABA without preincubation.

[³H]Ro 15-4513-binding assay to recombinant GABA_ARs. HEK293 cells were maintained in DMEM supplemented with 10% fetal bovine serum and seeded to a density of 500,000 cells on poly-lysine-coated 100-mm culture dishes 1 day before transfection. Cells were transfected with plasmids containing the subunit combination $\alpha 1\beta 2\gamma 2$ or $\alpha 1(H101R)\beta 2\gamma 2$ (7 mg total DNA, ratio 1:1:2) using jetPEI transfection reagent (Polyplus transfection). Twenty-four hours after transfection, HEK293 cells were harvested in PBS. HEK293 cells were homogenized in 10 mM Tris pH 7.5, protease inhibitor cocktail (complete Mini, Roche Applied Science). Aliquots of the homogenate prepared from HEK293 cells expressing the $\alpha 1\beta 2\gamma 2$ or $\alpha 1(H101R)\beta 2\gamma 2$ subunit combination were incubated with increasing concentrations of MDZ and 6.3 nM [³H]Ro15-4513 (22.7 Ci mmol⁻¹, PerkinElmer) in a total volume of 200 μ l for 90 min on ice. Subsequently, the samples were filtered on glass fibre filters using a 12-channel semiautomated cell harvester (Scatron) and washed with ice-cold buffer (10 mM Tris-HCl pH 7.4). Nonspecific [³H]Ro15-4513 binding was measured using 10 μ M flumazenil. The radioactivity of the filters was determined by liquid scintillation counting using a Tricarb 2500 liquid scintillation analyser.

Receptor occupancy. Mice were killed immediately after completion of the behavioural tests, that is, 90 min after DZP administration and 60 min after MDZ administration. Brains and spinal cords were removed and kept frozen until further processing. Percentage of receptors occupied by DZP or MDZ was determined *ex vivo* with [³H]flumazenil as radioligand⁴⁷. Frozen brains or lumbar spinal cords were rapidly homogenized in 20 volumes 10 mM Tris pH 7.4, 100 mM KCl and 100 μ l aliquots were immediately added to 400 μ l ice-cold buffer containing 4 nM [³H]flumazenil. After 30 s of incubation, samples were filtered on Whatman GF/C glass fibre filters and washed three times with 4 ml ice-cold buffer. Radioactivity retained on the filters was determined by scintillation counting. To determine nonspecific [³H]flumazenil binding, mice were used that had been injected i.p. with 5 mg kg⁻¹ bretazenil (dissolved in PEG300) to fully occupy all BDZ-binding sites (100% receptor occupancy⁴⁷).

References

- Zeilhofer, H. U., Benke, D. & Yévenes, G. E. Chronic pain states: pharmacological strategies to restore diminished inhibitory spinal pain control. *Annu. Rev. Pharmacol. Toxicol.* **52**, 111–133 (2012).
- De Koninck, Y. Altered chloride homeostasis in neurological disorders: a new target. *Curr. Opin. Pharmacol.* **7**, 93–99 (2007).
- Kuner, R. Central mechanisms of pathological pain. *Nat. Med.* **16**, 1258–1266 (2010).
- Sandkühler, J. Models and mechanisms of hyperalgesia and allodynia. *Physiol. Rev.* **89**, 707–758 (2009).
- Knabl, J. *et al.* Reversal of pathological pain through specific spinal GABA_A receptor subtypes. *Nature* **451**, 330–334 (2008).
- Sigel, E., Baur, R., Boulineau, N. & Minier, F. Impact of subunit positioning on GABA_A receptor function. *Biochem. Soc. Trans.* **34**, 868–871 (2006).
- Wieland, H. A., Luddens, H. & Seeburg, P. H. A single histidine in GABA_A receptors is essential for benzodiazepine agonist binding. *J. Biol. Chem.* **267**, 1426–1429 (1992).
- Benson, J. A., Löw, K., Keist, R., Möhler, H. & Rudolph, U. Pharmacology of recombinant γ -aminobutyric acid_A receptors rendered diazepam-insensitive by point-mutated α -subunits. *FEBS Lett.* **431**, 400–404 (1998).
- Möhler, H., Fritschy, J. M., Vogt, K., Crestani, F. & Rudolph, U. Pathophysiology and pharmacology of GABA_A receptors. *Handb. Exp. Pharmacol.* 225–247 (2005).
- McKernan, R. M. *et al.* Sedative but not anxiolytic properties of benzodiazepines are mediated by the GABA_A receptor $\alpha 1$ subtype. *Nat. Neurosci.* **3**, 587–592 (2000).
- Rudolph, U. *et al.* Benzodiazepine actions mediated by specific γ -aminobutyric acid_A receptor subtypes. *Nature* **401**, 796–800 (1999).
- Löw, K. *et al.* Molecular and neuronal substrate for the selective attenuation of anxiety. *Science* **290**, 131–134 (2000).
- Rudolph, U. & Knöflach, F. Beyond classical benzodiazepines: novel therapeutic potential of GABA_A receptor subtypes. *Nat. Rev. Drug Discov.* **10**, 685–697 (2011).
- Araujo, F. *et al.* Molecular and pharmacological characterization of native cortical γ -aminobutyric acid_A receptors containing both $\alpha 1$ and $\alpha 3$ subunits. *J. Biol. Chem.* **271**, 27902–27911 (1996).
- Benke, D. *et al.* Analysis of the presence and abundance of GABA_A receptors containing two different types of α subunits in murine brain using point-mutated α subunits. *J. Biol. Chem.* **279**, 43654–43660 (2004).
- Balic, E., Rudolph, U., Fritschy, J. M., Möhler, H. & Benke, D. The $\alpha 5(H105R)$ mutation impairs $\alpha 5$ selective binding properties by altered positioning of the $\alpha 5$ subunit in GABA_A receptors containing two distinct types of α subunits. *J. Neurochem.* **110**, 244–254 (2009).
- Bennett, G. J. & Xie, Y. K. A peripheral mononeuropathy in rat that produces disorders of pain sensation like those seen in man. *Pain* **33**, 87–107 (1988).
- Rivas, F. M. *et al.* Antiseizure activity of novel γ -aminobutyric acid_A receptor subtype-selective benzodiazepine analogues in mice and rat models. *J. Med. Chem.* **52**, 1795–1798 (2009).
- Fischer, B. D. *et al.* Anxiolytic-like effects of 8-acetylene imidazobenzodiazepines in a rhesus monkey conflict procedure. *Neuropharmacology* **59**, 612–618 (2010).
- Di Lio, A. *et al.* HZ166, a novel GABA_A receptor subtype-selective benzodiazepine site ligand, is antihyperalgesic in mouse models of inflammatory and neuropathic pain. *Neuropharmacology* **60**, 626–632 (2011).
- Drexler, B., Zinser, S., Hentschke, H. & Antkowiak, B. Diazepam decreases action potential firing of neocortical neurons via two distinct mechanisms. *Anesth. Analg.* **111**, 1394–1399 (2010).
- Walters, R. J., Hadley, S. H., Morris, K. D. & Amin, J. Benzodiazepines act on GABA_A receptors via two distinct and separable mechanisms. *Nat. Neurosci.* **3**, 1274–1281 (2000).
- Crestani, F. *et al.* Molecular targets for the myorelaxant action of diazepam. *Mol. Pharmacol.* **59**, 442–445 (2001).
- Dias, R. *et al.* Evidence for a significant role of $\alpha 3$ -containing GABA_A receptors in mediating the anxiolytic effects of benzodiazepines. *J. Neurosci.* **25**, 10682–10688 (2005).
- Jacob, T. C. *et al.* Benzodiazepine treatment induces subtype-specific changes in GABA_A receptor trafficking and decreases synaptic inhibition. *Proc. Natl Acad. Sci. USA* **109**, 18595–18600 (2012).
- Knabl, J., Zeilhofer, U. B., Crestani, F., Rudolph, U. & Zeilhofer, H. U. Genuine antihyperalgesia by systemic diazepam revealed by experiments in GABA_A receptor point-mutated mice. *Pain* **141**, 233–238 (2009).
- Hofmann, M. *et al.* Assessment of the effects of NS11394 and L-838417, $\alpha 2/3$ subunit-selective GABA_A receptor-positive allosteric modulators, in tests for pain, anxiety, memory and motor function. *Behav. Pharmacol.* **23**, 790–801 (2012).
- Nickolls, S. *et al.* A comparison of the $\alpha 2/3/5$ selective positive allosteric modulators L-838,417 and TPA023 in preclinical models of inflammatory and neuropathic pain. *Adv. Pharmacol. Sci.* **2011**, 608912 (2011).
- Munro, G. *et al.* Comparison of the novel subtype-selective GABA_A receptor-positive allosteric modulator NS11394 [3'-[5-(1-hydroxy-1-methyl-ethyl)-benzimidazol-1-yl]-biphenyl-2-carbonitrile] with diazepam, zolpidem,

- bretazenil, and gaboxadol in rat models of inflammatory and neuropathic pain. *J. Pharmacol. Exp. Ther.* **327**, 969–981 (2008).
30. Richards, B. L., Whittle, S. L. & Buchbinder, R. Muscle relaxants for pain management in rheumatoid arthritis. *Cochrane Database Syst. Rev.* **1**, CD008922 (2012).
 31. Corrigan, R., Derry, S., Wiffen, P. J. & Moore, R. A. Clonazepam for neuropathic pain and fibromyalgia in adults. *Cochrane Database Syst. Rev.* **5**, CD009486 (2012).
 32. Tan, K. R. *et al.* Neural bases for addictive properties of benzodiazepines. *Nature* **463**, 769–774 (2010).
 33. Fritschy, J. M. & Möhler, H. GABA_A-receptor heterogeneity in the adult rat brain: differential regional and cellular distribution of seven major subunits. *J. Comp. Neurol.* **359**, 154–194 (1995).
 34. Bohlhalter, S., Weinmann, O., Möhler, H. & Fritschy, J. M. Laminar compartmentalization of GABA_A-receptor subtypes in the spinal cord: an immunohistochemical study. *J. Neurosci.* **16**, 283–297 (1996).
 35. Engin, E. *et al.* Neural basis of benzodiazepine reward: requirement for $\alpha 2$ containing GABA_A receptors in the nucleus accumbens. *Neuropsychopharmacology* **39**, 1805–1815 (2014).
 36. Reynolds, L. M. *et al.* Differential roles of GABA_A receptor subtypes in benzodiazepine-induced enhancement of brain-stimulation reward. *Neuropsychopharmacology* **37**, 2531–2540 (2012).
 37. Tatsuo, M. A., Salgado, J. V., Yokoro, C. M., Duarte, I. D. & Francischi, J. N. Midazolam-induced hyperalgesia in rats: modulation via GABA_A receptors at supraspinal level. *Eur. J. Pharmacol.* **370**, 9–15 (1999).
 38. Luger, T. J., Hayashi, T., Weiss, C. G. & Hill, H. F. The spinal potentiating effect and the supraspinal inhibitory effect of midazolam on opioid-induced analgesia in rats. *Eur. J. Pharmacol.* **275**, 153–162 (1995).
 39. Harris, J. A. & Westbrook, R. F. Effects of benzodiazepine microinjection into the amygdala or periaqueductal gray on the expression of conditioned fear and hypoalgesia in rats. *Behav. Neurosci.* **109**, 295–304 (1995).
 40. Vuilleumier, P. H., Besson, M., Desmeules, J., Arendt-Nielsen, L. & Curatolo, M. Evaluation of anti-hyperalgesic and analgesic effects of two benzodiazepines in human experimental pain: a randomized placebo-controlled study. *PLoS ONE* **8**, e43896 (2013).
 41. Atack, J. *et al.* Preclinical and clinical pharmacology of TPA023B, a GABA_A receptor $\alpha 2/\alpha 3$ subtype-selective partial agonist. *J. Psychopharmacol.* **25**, 329–344 (2011).
 42. Dawson, G. R. *et al.* An inverse agonist selective for $\alpha 5$ subunit-containing GABA_A receptors enhances cognition. *J. Pharmacol. Exp. Ther.* **316**, 1335–1345 (2006).
 43. Rudolph, U. & Möhler, H. GABA_A receptor subtypes: therapeutic potential in Down syndrome, affective disorders, schizophrenia, and autism. *Annu. Rev. Pharmacol. Toxicol.* **54**, 483–507 (2014).
 44. Crestani, F. *et al.* Trace fear conditioning involves hippocampal $\alpha 5$ GABA_A receptors. *Proc. Natl Acad. Sci. USA* **99**, 8980–8985 (2002).
 45. Paul, J., Zeilhofer, H. U. & Fritschy, J. M. Selective distribution of GABA_A receptor subtypes in mouse spinal dorsal horn neurons and primary afferents. *J. Comp. Neurol.* **520**, 3895–3911 (2012).
 46. Paul, J. *et al.* Antihyperalgesia by $\alpha 2$ -GABA_A receptors occurs via a genuine spinal action and does not involve supraspinal sites. *Neuropsychopharmacology* **39**, 477–487 (2014).
 47. Li, J., Fish, R. L., Cook, S. M., Tattersall, F. D. & Atack, J. R. Comparison of *in vivo* and *ex vivo* [³H]flumazenil binding assays to determine occupancy at the benzodiazepine binding site of rat brain GABA_A receptors. *Neuropharmacology* **51**, 168–172 (2006).

Acknowledgements

We are grateful to Dr James M. Cook and Michael M. Poe, University of Wisconsin, Milwaukee, for their generous gift of HZ-166, to Dr Jean-Marc Fritschy for help with the immunocytochemistry, to Dr Hendrik Wildner for critical reading of the manuscript, to Isabelle Kellenberger for mouse genotyping and to Carmen Brazerol, Conny Leuzinger and Dennis Kwame for animal care taking. This work was in part supported by grants from the Swiss National Science Foundation to H.U.Z. (131093 and 156393) and an Advanced Investigator ERC grant to HUZ (DHIS, 250128). U.R. is supported by grants from the National Institutes of Health (R03MH094834 and R03DA033491).

Author contributions

W.T.R. performed all experiments, except the electrophysiological measurement, and analysed the data. D.B. performed and analysed the radio ligand-binding experiments. M.A.A. performed and analysed the electrophysiology experiments. U.R. helped design the experiments. H.U.Z. designed experiments, analysed data and wrote the manuscript. All authors made comments on the manuscript.

Additional information

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Ralvenius, W. T. *et al.* Analgesia and unwanted benzodiazepine effects in point-mutated mice expressing only one benzodiazepine-sensitive GABA_A receptor subtype. *Nat. Commun.* **6**:6803 doi: 10.1038/ncomms7803 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>